arXiv:1105.0093v1 [math.NT] 30 Apr 2011

Some new identities on the twisted (h, q)-Euler numbers and q-Bernstein polynomials

D. V. Dolgy[1], D. J. Kang[2], T. Kim[3], and B. Lee[4]

[1]Hanrimwon, Kwangwoon University, Seoul 139-701, Korea

[2] Information Technology Service,
Kyungpook National University, Taegu 702-701, Korea

[3]Division of General Education-Mathematics,
Kwangwoon University, Seoul 139-701, Korea

[4]Department of Wireless Communications Engineering,
Kwangwoon University, Seoul 139-701, Korea

Abstract In this paper we give some interesting relationship between the q-Bernstein polynomials and the twisted (h, q)-Euler numbers by using fermionic p-adic q-integral on $\mathbb{Z}_p$.

2000 Mathematics Subject Classification - 11B68, 11S40, 11S80

Key words- Euler numbers and polynomials, twisted (h, q)-Euler numbers and polynomials, Bernstein polynomials, q-Bernstein polynomials

1. Introduction

Let p be a fixed odd prime number. Throughout this paper, we always make use of the following notations: $\mathbb{Z}$ denotes the ring of rational integers, $\mathbb{Z}_p$ denotes the ring of p-adic rational integers, $\mathbb{Q}_p$ denotes the field of p-adic rational numbers, and $\mathbb{C}_p$ denotes the completion of algebraic closure of $\mathbb{Q}_p$, respectively. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{Z}_+ = \mathbb{N} \cup \{0\}$. Let $C_{p^n} = \{\zeta | \zeta^{p^n} = 1\}$ be the cyclic group of order p^n and let

$$T_p = \lim_{n\to\infty} C_{p^n} = \cup_{n\geq 1} C_{p^n}.$$

The p-adic absolute value is defined by $|x|_p = \frac{1}{p^r}$, where $x = p^r \frac{s}{t}$($r \in \mathbb{Q}$ and $s, t \in \mathbb{Z}$ with $(s,t) = (p,s) = (p,t) = 1)$. In this paper we assume that $q \in \mathbb{C}_p$ with $|q-1|_p < 1$ as an indeterminate. The q-number is defined by

$$[x]_q = \frac{1-q^x}{1-q}, \text{ see [1-18]}.$$

Note that $\lim_{q\to 1}[x]_q = x$. For

$$f \in UD(\mathbb{Z}_p) = \{f | f : \mathbb{Z}_p \to \mathbb{C}_p \text{ is uniformly differentiable function}\},$$

the fermionic p-adic q-integral on $\mathbb{Z}_p$ is defined as

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N\to\infty} \frac{1+q}{1+q^{p^N}} \sum_{x=0}^{p^N-1} f(x)(-q)^x, \text{ see [1-5] }. \tag{1}$$

From (1), we note that

$$q^n I_{-q}(f_n) = (-1)^n I_{-q}(f) + [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} q^l f(l),$$

where $f_n(x) = f(x+n)$ for $n \in \mathbb{N}$. For $k, n \in \mathbb{Z}_+$ and $x \in [0,1]$, Kim defined q-Bernstein polynomials, which are different q-Bernstein polynomials of Phillips, as follows:

$$B_{k,n}(x,q) = \binom{n}{k} [x]_q^k [1-x]_{q^{-1}}^{n-k}, \text{ see [4]}. \tag{2}$$

In [9], the p-adic extension of (2) is given by

$$B_{k,n}(x,q) = \binom{n}{k} [x]_q^k [1-x]_{q^{-1}}^{n-k}, \text{ where } x \in \mathbb{Z}_p, \text{ and } n, k \in \mathbb{Z}_+. \tag{3}$$

For $h \in \mathbb{Z}$ and $\zeta \in T_p$, let us consider the twisted (h,q)-Euler polynomials as follows:

$$E_{n,q,\zeta}^{(h)}(x) = \int_{\mathbb{Z}_p} [x+y]_q^n \zeta^y q^{(h-1)y} d\mu_{-q}(y), \text{ for } n \in \mathbb{Z}_+. \tag{4}$$

In the special case, $x = 0$, $E_{n,q,\zeta}^{(h)}(0) = E_{n,q,\zeta}^{(h)}$ are called the n-th twisted (h,q)-Euler numbers.

In this paper we investigate some relations between the q-Bernstein polynomials and the twisted (h,q)-Euler numbers. From these relations, we derive some interesting identities on the the twisted (h,q)-Euler numbers and polynomials.

2. On the twisted (h,q)-Euler numbers and polynomials

From (4), we note that

$$\begin{aligned} E_{n,q,\zeta}^{(h)}(x) &= \int_{\mathbb{Z}_p} [x+y]_q^n \zeta^y q^{(h-1)y} d\mu_{-q}(y) \\ &= \frac{[2]_q}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} (-1)^l q^{lx} \left(\frac{1}{1+q^{h+l}\zeta} \right) \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m \zeta^m q^{hm} [x+m]_q^n, \end{aligned}$$

and

$$\begin{aligned} E_{n,q,\zeta}^{(h)}(x) &= \int_{\mathbb{Z}_p} [x+y]_q^n \zeta^y q^{(h-1)y} d\mu_{-q}(y) \\ &= \sum_{l=0}^{n} \binom{n}{l} [x]_q^{n-l} q^{lx} \int_{\mathbb{Z}_p} [y]_q^l \zeta^y q^{(h-1)y} d\mu_{-q}(y) \qquad (5) \\ &= \sum_{l=0}^{n} \binom{n}{l} [x]_q^{n-l} q^{lx} E_{l,q,\zeta}^{(h)}. \end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 1. For $n \in \mathbb{Z}_+$ and $\zeta \in T_p$, we have

$$E_{n,q,\zeta}^{(h)}(x) = [2]_q \sum_{m=0}^{\infty} (-1)^m \zeta^m q^{hm} [x+m]_q^n.$$

Furthermore,

$$\begin{aligned} E_{n,q,\zeta}^{(h)}(x) &= \sum_{l=0}^{n} \binom{n}{l} [x]_q^{n-l} q^{lx} E_{l,q,\zeta}^{(h)} \\ &= ([x]_q + q^x E_{q,\zeta}^{(h)})^n, \end{aligned}$$

with usual convention about replacing $(E_{q,\zeta}^{(h)})^n$ by $E_{n,q,\zeta}^{(h)}$.

Let

$$F_{q,\zeta}^{(h)}(t,x) = \sum_{n=0}^{\infty} E_{n,q,\zeta}^{(h)}(x) \frac{t^n}{n!}.$$

Then we see that

$$F_{q,\zeta}^{(h)}(t,x) = [2]_q \sum_{m=0}^{\infty} (-1)^m \zeta^m q^{mh} e^{[x+m]_q t}. \qquad (6)$$

In the special case, $x=0$, let $F_{q,\zeta}^{(h)}(t,0) = F_{q,\zeta}^{(h)}(t)$.

By (4), we get

$$
\begin{aligned}
E^{(h)}_{n,q^{-1},\zeta^{-1}}(1-x) &= \int_{\mathbb{Z}_p} [1-x+y]^n_{q^{-1}} \zeta^{-y} q^{-(h-1)y} d\mu_{-q^{-1}}(y) \\
&= \frac{[2]_q}{(1-q^{-1})^n} \sum_{l=0}^{n} \binom{n}{l} (-1)^l q^{h-1} \zeta \left(\frac{q^{lx}}{1+q^{h+l}\zeta} \right) \\
&= (-1)^n \zeta q^{n+h-1} \left(\frac{[2]_q}{(1-q)^n} \sum_{l=0}^{n} \binom{n}{l} (-1)^l \frac{q^{lx}}{1+q^{h+l}\zeta} \right) \\
&= (-1)^n \zeta q^{n+h-1} E^{(h)}_{n,q,\zeta}(x).
\end{aligned}
$$

Therefore, we obtain the following theorem.

Theorem 2. Let $F^{(h)}_{q,\zeta}(t,x) = \sum_{n=0}^{\infty} E^{(h)}_{n,q,\zeta}(x) \frac{t^n}{n!}$. Then we have

$$F^{(h)}_{q^{-1},\zeta^{-1}}(t,1-x) = \zeta q^{h-1} F^{(h)}_{q,\zeta}(-qt,x).$$

Moreover,

$$E^{(h)}_{n,q^{-1},\zeta^{-1}}(1-x) = (-1)^n \zeta q^{n+h-1} E^{(h)}_{n,q,\zeta}(x) \text{ for } n \in \mathbb{Z}_+.$$

From (6), we note that

$$q^h \zeta F^{(h)}_{q,\zeta}(t,1) + F^{(h)}_{q,\zeta}(t) = [2]_q. \tag{7}$$

By (7), we get the following recurrence formula:

$$E^{(h)}_{0,q,\zeta} = \frac{1+q}{1+q^h\zeta}, \text{ and } q^h \zeta E^{(h)}_{n,q,\zeta}(1) + E^{(h)}_{n,q,\zeta} = 0 \text{ if } n > 0. \tag{8}$$

By (8) and Theorem 1, we obtain the following theorem.

Theorem 3. For $n \in \mathbb{Z}_+$ and $\zeta \in T_p$, we have

$$E^{(h)}_{0,q,\zeta} = \frac{1+q}{1+q^h\zeta}, \text{ and } q^h \zeta (qE^{(h)}_{q,\zeta} + 1)^n + E^{(h)}_{n,q,\zeta} = 0 \text{ if } n > 0,$$

with usual convention about replacing $(E^{(h)}_{q,\zeta})^n$ by $E^{(h)}_{n,q,\zeta}$.

From Theorem 3, we note that

$$\begin{aligned}
&q^{2h}\zeta^2 E^{(h)}_{n,q,\zeta}(2) - \frac{1+q}{1+q^h\zeta}q^{2h}\zeta^2 - \frac{1+q}{1+q^h\zeta}q^h\zeta \\
&= q^{2h}\zeta^2 \sum_{l=0}^{n}\binom{n}{l}q^l(qE^{(h)}_{q,\zeta}+1)^l - \frac{1+q}{1+q^h\zeta}q^{2h}\zeta^2 - \frac{1+q}{1+q^h\zeta}q^h\zeta \\
&= q^{2h}\zeta^2 \sum_{l=1}^{n}\binom{n}{l}q^l(qE^{(h)}_{q,\zeta}+1)^l - \frac{1+q}{1+q^h\zeta}q^h\zeta \\
&= -q^h\zeta \sum_{l=1}^{n}\binom{n}{l}q^l E^{(h)}_{l,q,\zeta} - \frac{1+q}{1+q^h\zeta}q^h\zeta \\
&= -q^h\zeta \sum_{l=0}^{n}\binom{n}{l}q^l E^{(h)}_{l,q,\zeta} \\
&= -q^h\zeta E^{(h)}_{n,q,\zeta}(1) = E^{(h)}_{n,q,\zeta} \text{ if } n > 0.
\end{aligned}$$

Therefore, we obtain the following theorem.

Theorem 4. For $n \in \mathbb{N}$, we have

$$q^{2h}\zeta^2 E^{(h)}_{n,q,\zeta}(2) = E^{(h)}_{n,q,\zeta} + \frac{1+q}{1+q^h\zeta}q^{2h}\zeta^2 + \frac{1+q}{1+q^h\zeta}q^h\zeta.$$

By Theorem 2, we see that

$$\begin{aligned}
&q^{h-1}\zeta \int_{\mathbb{Z}_p} [1-x]^n_{q^{-1}} q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\
&= (-1)^n q^{n+h-1}\zeta \int_{\mathbb{Z}_p} [x-1]^n_q q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\
&= (-1)^n q^{n+h-1}\zeta E^{(h)}_{n,q,\zeta}(-1) = E^{(h)}_{n,q^{-1},\zeta^{-1}}(2).
\end{aligned} \tag{9}$$

Therefore, we obtain the following theorem.

Theorem 5. For $n \in \mathbb{Z}_+$, we have

$$q^{h-1}\zeta \int_{\mathbb{Z}_p} [1-x]^n_{q^{-1}} q^{(h-1)x}\zeta^x d\mu_{-q}(x) = E^{(h)}_{n,q^{-1},\zeta^{-1}}(2).$$

Let $n \in \mathbb{N}$. By Theorem 5 and Theorem 4, we get

$$\begin{aligned}
&q^{h-1}\zeta \int_{\mathbb{Z}_p} [1-x]^n_{q^{-1}} q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\
&= q^{2h}\zeta^2 E^{(h)}_{n,q^{-1},\zeta^{-1}} + q^{h-1}\zeta\left(\frac{1+q}{1+q^h\zeta}\right) + q^{2h-1}\zeta^2\left(\frac{1+q}{1+q^h\zeta}\right).
\end{aligned} \tag{10}$$

From (10), we have

$$\int_{\mathbb{Z}_p}[1-x]_{q^{-1}}^n q^{(h-1)x}\zeta^x d\mu_{-q}(x)=q^{h+1}\zeta E_{n,q^{-1},\zeta^{-1}}^{(h)}+\left(\frac{1+q}{1+q^h\zeta}\right)+q^h\zeta\left(\frac{1+q}{1+q^h\zeta}\right).$$

Therefore, we obtain the following corollary.

Corollary 6. For $n\in\mathbb{N}$, we have

$$\int_{\mathbb{Z}_p}[1-x]_{q^{-1}}^n q^{(h-1)x}\zeta^x d\mu_{-q}(x)=q^{h+1}\zeta E_{n,q^{-1},\zeta^{-1}}^{(h)}+[2]_q.$$

For $x\in\mathbb{Z}_p$, the p-adic analogues of q-Bernstein polynomials are given by

$$B_{k,n}(x,q)=\binom{n}{k}[x]_q^k[1-x]_{q^{-1}}^{n-k},\ \text{where } n,k\in\mathbb{Z}_+. \tag{11}$$

By (11), we get the symmetry of q-Bernstein polynomials as follows:

$$B_{k,n}(x,q)=B_{n-k,n}(1-x,q^{-1}),\ \text{see [7]}. \tag{12}$$

Thus, by Corollary 6, (11), and (12), we see that

$$\begin{aligned}\int_{\mathbb{Z}_p}B_{k,n}(x,q)q^{(h-1)x}\zeta^x d\mu_{-q}(x)&=\int_{\mathbb{Z}_p}B_{n-k,n}(1-x,q^{-1})q^{(h-1)x}\zeta^x d\mu_{-q}(x)\\&=\binom{n}{k}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k+l}\int_{\mathbb{Z}_p}[1-x]_{q^{-1}}^{n-l}q^{(h-1)x}\zeta^x d\mu_{-q}(x)\\&=\binom{n}{k}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k+l}\left(q^{h+1}\zeta E_{n-l,q^{-1},\zeta^{-1}}^{(h)}+[2]_q\right).\end{aligned}$$

For $n,k\in\mathbb{Z}_+$ with $n>k$, we have

$$\begin{aligned}&\int_{\mathbb{Z}_p}B_{k,n}(x,q)q^{(h-1)x}\zeta^x d\mu_{-q}(x)\\&=\binom{n}{k}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k+l}\left(q^{h+1}\zeta E_{n,q^{-1},\zeta^{-1}}^{(h)}+[2]_q\right)\\&=\begin{cases}q^{h+1}\zeta E_{n,q^{-1},\zeta^{-1}}^{(h)}+[2]_q, & \text{if } k=0,\\ q^{h+1}\zeta\binom{n}{k}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k+l}q^{h+1}\zeta E_{n-l,q^{-1},\zeta^{-1}}^{(h)}, & \text{if } k>0.\end{cases}\end{aligned} \tag{13}$$

Let us take the fermionic q-integral on $\mathbb{Z}_p$ for the q-Bernstein polynomials of degree n as follows:

$$
\begin{aligned}
\int_{\mathbb{Z}_p} B_{k,n}(x,q) q^{(h-1)x}\zeta^x d\mu_{-q}(x) &= \binom{n}{k}\int_{\mathbb{Z}_p} [x]_q^k [1-x]_{q^{-1}}^{n-k} q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\
&= \binom{n}{k}\sum_{l=0}^{n-k}\binom{n-k}{l}(-1)^l E_{l+k,q,\zeta}^{(h)}.
\end{aligned} \tag{14}
$$

Therefore, by (13) and (14), we obtain the following theorem.

Theorem 7. Let $n, k \in \mathbb{Z}_+$ with $n > k$. Then we have

$$
\begin{aligned}
&\int_{\mathbb{Z}_p} B_{k,n}(x,q) q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\
&= \begin{cases} q^{h+1}\zeta E_{n,q^{-1},\zeta^{-1}}^{(h)} + [2]_q, & \text{if } k = 0, \\ q^{h+1}\zeta \binom{n}{k}\sum_{l=0}^{k}\binom{k}{l}(-1)^{k+l} E_{n-l,q^{-1},\zeta^{-1}}^{(h)}, & \text{if } k > 0. \end{cases}
\end{aligned}
$$

Moreover,

$$
\begin{aligned}
&\sum_{l=0}^{n-k}\binom{n-k}{l}(-1)^l E_{l+k,q,\zeta}^{(h)} \\
&= \begin{cases} q^{h+1}\zeta E_{n,q^{-1},\zeta^{-1}}^{(h)} + [2]_q, & \text{if } k = 0, \\ q^{h+1}\zeta \sum_{l=0}^{k}\binom{k}{l}(-1)^{k+l} E_{n-l,q^{-1},\zeta^{-1}}^{(h)}, & \text{if } k > 0. \end{cases}
\end{aligned}
$$

Let $n_1, n_2, k \in \mathbb{Z}_+$ with $n_1 + n_2 > 2k$. Then we get

$$
\begin{aligned}
&\int_{\mathbb{Z}_p} B_{k,n_1}(x,q) B_{k,n_2}(x,q) q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\
&= \binom{n_1}{k}\binom{n_2}{k}\sum_{l=0}^{2k}\binom{2k}{l}(-1)^{l+2k}\int_{\mathbb{Z}_p}[1-x]_{q^{-1}}^{n_1+n_2-l} q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\
&= \binom{n_1}{k}\binom{n_2}{k}\sum_{l=0}^{2k}\binom{2k}{l}(-1)^{l+2k}\left(q^{h+1}\zeta E_{n_1+n_2-l,q^{-1},\zeta^{-1}}^{(h)} + [2]_q\right) \\
&= \begin{cases} q^{h+1}\zeta E_{n_1+n_2,q^{-1},\zeta^{-1}}^{(h)} + [2]_q, & \text{if } k = 0, \\ \binom{n_1}{k}\binom{n_2}{k}\sum_{l=0}^{2k}\binom{2k}{l}(-1)^{2k+l}\left(q^{h+1}\zeta E_{n_1+n_2-l,q^{-1},\zeta^{-1}}^{(h)} + [2]_q\right), & \text{if } k \neq 0. \end{cases}
\end{aligned} \tag{15}
$$

Therefore, by (15), we obtain the following theorem.

Theorem 8. For $n_1, n_2, k \in \mathbb{Z}_+$ with $n_1 + n_2 > 2k$, we have

$$\begin{aligned} &\int_{\mathbb{Z}_p} B_{k,n_1}(x,q) B_{k,n_2}(x,q) q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\ &= \begin{cases} q^{h+1}\zeta E^{(h)}_{n_1+n_2,q^{-1},\zeta^{-1}} + [2]_q, & \text{if } k = 0, \\ q^{h+1}\zeta \binom{n_1}{k}\binom{n_2}{k} \sum_{l=0}^{2k}\binom{2k}{l}(-1)^{2k+l} E^{(h)}_{n_1+n_2-l,q^{-1},\zeta^{-1}}, & \text{if } k \neq 0. \end{cases} \end{aligned}$$

From the binomial theorem, we can derive the following equation.

$$\begin{aligned} &\int_{\mathbb{Z}_p} B_{k,n_1}(x,q) B_{k,n_2}(x,q) q^{(h-1)x}\zeta^x d\mu_{-q}(x) \\ &= \binom{n_1}{k}\binom{n_2}{k} \sum_{l=0}^{n_1+n_2-2k} (-1)^l \binom{n_1+n_2-2k}{l} \int_{\mathbb{Z}_p} [x]_q^{2k+l} q^{(h-1)x}\zeta^x d\mu_{-q}(x) \qquad (16) \\ &= \binom{n_1}{k}\binom{n_2}{k} \sum_{l=0}^{n_1+n_2-2k} (-1)^l \binom{n_1+n_2-2k}{l} E^{(h)}_{2k+l,q,\zeta}. \end{aligned}$$

Thus, by Theorem 8 and (16), we obtain the following corollary.

Corollary 9. Let $n_1, n_2, k \in \mathbb{Z}_+$ with $n_1 + n_2 > 2k$. Then we have

$$\begin{aligned} &\sum_{l=0}^{n_1+n_2-2k} (-1)^l \binom{n_1+n_2-2k}{l} E^{(h)}_{2k+l,q} \\ &= \begin{cases} q^{h+1}\zeta E^{(h)}_{n_1+n_2,q^{-1},\zeta^{-1}} + [2]_q, & \text{if } k = 0, \\ q^{h+1}\zeta \sum_{l=0}^{2k}\binom{2k}{l}(-1)^{2k+l} E^{(h)}_{n_1+n_2-l,q^{-1},\zeta^{-1}}, & \text{if } k > 0. \end{cases} \end{aligned}$$

For $x \in \mathbb{Z}_p$ and $s \in \mathbb{N}$ with $s \geq 2$, let $n_1, n_2, \ldots, n_s, k \in \mathbb{Z}_+$ with $n_1 + \cdots + n_s > sk$. Then we take the fermionic p-adic q-integral on $\mathbb{Z}_p$ for the q-Bernstein polynomials of degree n as follows:

$$\begin{aligned}
&\int_{\mathbb{Z}_p} \underbrace{B_{k,n_1}(x,q)\cdots B_{k,n_s}(x,q)}_{s-\text{times}} q^{(h-1)x}\zeta^x d\mu_{-q}(x)\\
&= \binom{n_1}{k}\cdots\binom{n_s}{k}\int_{\mathbb{Z}_p}[x]_q^{sk}[1-x]_{q^{-1}}^{n_1+\cdots+n_s-sk}q^{(h-1)x}\zeta^x d\mu_{-q}(x)\\
&= \binom{n_1}{k}\cdots\binom{n_s}{k}\sum_{l=0}^{sk}\binom{sk}{l}(-1)^{l+sk}\int_{\mathbb{Z}_p}[1-x]_{q^{-1}}^{n_1+\cdots+n_s-l}q^{(h-1)x}\zeta^x d\mu_{-q}(x)\\
&= \binom{n_1}{k}\cdots\binom{n_s}{k}\sum_{l=0}^{sk}\binom{sk}{l}(-1)^{l+sk}\left(q^{h+1}\zeta E^{(h)}_{n_1+\cdots+n_s-l,q^{-1},\zeta^{-1}}+[2]_q\right)\\
&= \begin{cases} q^{h+1}\zeta E^{(h)}_{n_1+\cdots+n_s,q^{-1},\zeta^{-1}}+[2]_q, & \text{if } k=0,\\[2ex] q^{h+1}\zeta\binom{n_1}{k}\cdots\binom{n_s}{k}\sum_{l=0}^{sk}\binom{sk}{l}(-1)^{l+sk}E^{(h)}_{n_1+\cdots+n_s-l,q^{-1},\zeta^{-1}}, & \text{if } k>0.\end{cases}
\end{aligned}\tag{17}$$

Therefore, by (17), we obtain the following theorem.

Theorem 10. For $s \in \mathbb{N}$ with $s \geq 2$, let $n_1, n_2, \ldots, n_s, k \in \mathbb{Z}_+$ with $n_1+\cdots+n_s > sk$. Then we get

$$\begin{aligned}
&\int_{\mathbb{Z}_p} \underbrace{B_{k,n_1}(x,q)\cdots B_{k,n_s}(x,q)}_{s-\text{times}} q^{(h-1)x}\zeta^x d\mu_{-q}(x)\\
&= \begin{cases} q^{h+1}\zeta E^{(h)}_{n_1+\cdots+n_s,q^{-1},\zeta^{-1}}+[2]_q, & \text{if } k=0,\\[2ex] q^{h+1}\zeta\binom{n_1}{k}\cdots\binom{n_s}{k}\sum_{l=0}^{sk}\binom{sk}{l}(-1)^{l+sk}E^{(h)}_{n_1+\cdots+n_s-l,q^{-1},\zeta^{-1}}, & \text{if } k>0.\end{cases}
\end{aligned}\tag{18}$$

By the definition of q-Bernstein polynomials and the binomial theorem, we easily get

$$\begin{aligned}
&\int_{\mathbb{Z}_p} \underbrace{B_{k,n_1}(x,q)\cdots B_{k,n_s}(x,q)}_{s-\text{times}} q^{(h-1)x}\zeta^x d\mu_{-q}(x)\\
&= \binom{n_1}{k}\cdots\binom{n_s}{k}\sum_{l=0}^{n_1+\cdots+n_s-sk}(-1)^l\binom{n_1+\cdots+n_s-sk}{l}\int_{\mathbb{Z}_p}[x]_q^{sk+l}q^{(h-1)x}\zeta^z d\mu_{-q}(x)\\
&= \binom{n_1}{k}\cdots\binom{n_s}{k}\sum_{l=0}^{n_1+\cdots+n_s-sk}(-1)^l\binom{n_1+\cdots+n_s-sk}{l}E^{(h)}_{sk+l,q,\zeta}.
\end{aligned}$$

Therefore, we have the following corollary.

Corollary 11. For $\zeta \in T_p, s \in \mathbb{N}$ with $s \geq 2$, let $n_1, n_2, \ldots, n_s, k \in \mathbb{Z}_+$ with $n_1 + \cdots + n_s > sk$. Then we have

$$\sum_{l=0}^{n_1+\cdots+n_s-sk} (-1)^l \binom{n_1 + \cdots + n_s - sk}{l} E^{(h)}_{sk+l,q,\zeta}$$
$$= \begin{cases} q^{h+1}\zeta E^{(h)}_{n_1+\cdots+n_s,q^{-1},\zeta^{-1}} + [2]_q, & \text{if } k = 0, \\ q^{h+1}\zeta \sum_{l=0}^{sk} \binom{sk}{l} (-1)^{l+sk} E^{(h)}_{n_1+\cdots+n_s-l,q^{-1},\zeta^{-1}}, & \text{if } k > 0. \end{cases}$$

References

[1] T. Kim, q-Euler numbers and polynomials associated with p-adic q-integrals, J. Nonlinear Math. Phys. 14(2007), 15-27.

[2] T. Kim, Some identities on the q-Euler polynomials of higher order and q-Stirling numbers by the fermionic p-adic integral on $\mathbb{Z}_p$, Russ. J. Math. Phys. 16(2009), 484-491.

[3] T. Kim, Barnes type multiple q-zeta function and q-Euler polynomials, J. Phys. A : Math. Theor. 43(2010) 255201, 11pp.

[4] T. Kim, A note on q-Bernstein polynomials, Russ. J. Math. Phys. 18(2011), 41-50.

[5] T. Kim, q-Volkenborn integration, Russ. J. Math. Phys. 9(2002), 288-299.

[6] T. Kim, q-Bernoulli numbers and polynomials associated with Gaussian binomial coefficients, Russ. J. Math. Phys. 15(2008), 51-57.

[7] T. Kim, J. Choi, Y. H. Kim, C. S. Ryoo, On the fermionic p-adic integral representation of Bernstein polynomials associated with Euler numbers and polynomials, J. Inequal. Appl. 2010(2010), Art ID 864247, 12pp.

[8] T. Kim, B. Lee, J. Choi, Y. H. Kim, A new approach of q-Euler numbers and polynomials, Proc. Jangjeon Math. Soc. 14(2011), 7-14.

[9] T. Kim, J. Choi, Y.-H. Kim, Some identities on the q-Bernstein polynomials, q-Stirling numbers and q-Bernoulli numbers, Adv. Stud. Contemp. Math. 20(2010), 335-341.

[10] T. Kim, Some identities for the Bernoulli, the Euler and the Genocchi numbers and polynomials, Adv. Stud. Contemp. Math. 20(2010), 23-28.

[11] T. Kim, The modified q-Euler numbers and polynomials, Adv. Stud. Contemp. Math. 16(2008), 161-170.

[12] T. Kim, Note on the Euler numbers and polynomials, Adv. Stud. Contemp. Math. 17(2008), 131-136.

[13] T. Kim, B. Lee, J. Choi, Y. H. Kim, S. H. Rim, On the q-Euler numbers and weighted q-Bernstein polynomials, Adv. Stud. Contemp. Matth. 21(2011), 13-18.

[14] S.-H. Rim, J.-H. Jin, E.-J. Moon, S.-J. Lee, On multiple interpolation function of the q-Genocchi polynomials, J. Inequal. Appl. 2010, Art ID 351419, 13pp.

[15] S.-H. Rim, S. J. Lee, E. J. Moon, J. H. Jin, On the q-Genocchi numbers and polynomials associated with q-zeta function, Proc. Jangjeon Math. Soc. 12(2009), 261-267.

[16] Y. Simsek, O. Yurekli, V. Kurt, On interpolation functions of the twisted generalized Frobinuous-Euler numbers, Adv. Stud. Contemp. Math. 14(2007), 49-68.

[17] Y. Simsek, Twisted p-adic (h,q)-L-functions, Comput. Math. Appl. 59(2010), 2097–2110 .

[18] H. Ozden, Y. Simsek, Interpolation function of the (h,q)-extension of twisted Euler numbers, Comput. Math. Appl. 56(2008), 898–908.